

Abstract Submitted  
for the DPP14 Meeting of  
The American Physical Society

**Experimental optimization of high magnetic Reynolds number, two-vortex flows on the Madison plasma dynamo experiment** DAVID WEISBERG, University of Wisconsin - Madison — Laminar counter-rotating two-vortex flows, predicted to excite a large-scale dynamo, are generated and optimized in the Madison plasma dynamo experiment (MPDX). Numerical simulations show that the topology of these simply-connected flows may be optimal for creating a plasma dynamo in the lab and predict a critical threshold of  $Rm_{crit} = \mu_0 \sigma LV = 250$  for optimal flows. MPDX plasmas are shown to exceed this critical  $Rm$ , generating large ( $L = 1.4$  m), hot ( $T_e > 10$  eV) plasmas where  $Rm = 600$ . Plasma flow is driven using eight thermally emissive LaB<sub>6</sub> cathodes which generate a  $J \times B$  torque at the magnetized edge of spherical He plasmas. Mach probe measurements show counter-rotating flows at speeds of  $V > 10$  km/s; the driven flow at the plasma edge viscously couples inward to the unmagnetized core via ion-ion collisions, and flow measurements along radial chords compare favorably to hydrodynamic calculations using Braginskii viscosity. To optimize flow for dynamo generation, cathode bias and positioning are varied along with plasma viscosity ( $\nu \sim T_i^{5/2}/n_i$ ) and the frictional neutral-ion drag force ( $\mu = L^2/(\nu\tau_{in})$ ). Prospects for observing a dynamo, hydrodynamic transitions to turbulence, and eventual large Rm fast dynamos will be presented.

David Weisberg  
University of Wisconsin - Madison

Date submitted: 27 Jun 2014

Electronic form version 1.4