

Abstract Submitted
for the DPP14 Meeting of
The American Physical Society

Recent High-Intensity Experiments at the Trident Laser¹ JAMES COBBLE, SASIKUMAR PALANIYAPPAN, CORT GAUTIER, YONGHO KIM, CHENGKUN HUANG, Los Alamos National Laboratory — With near-diffraction-limited irradiance of 2×10^{20} W/cm² on target and pre-pulse contrast better than 10^{-8} , we have accessed the regime of relativistic transparency (RT) at the Trident Laser. The goal was to assess electron debris emitted from the target rear surface with phase-contrast imaging (PCI) and current density measurements (hence, the total electron current). Companion diagnostics show whether the experiments are in the target-normal-sheath-acceleration mode or in the RT regime. The superb laser contrast allows us to shoot targets as thin as 50 nm. PCI at 527 nm is temporally resolved to 600 fs. It has shown the evolution of electron behavior over tens of ps, including thermal electrons accompanying the ion jet, accelerated to many tens of MeV earlier in time. Faraday-cup measurements indicate the transfer of many microC of charge during the laser drive. As a ride-along experiment using a gas Cherenkov detector (GCD), we have detected gamma rays of energy >5 MeV. This radiation has a prompt component and a lesser source, driven by accelerated ions, that is time resolved by the GCD. The ion time of flight is compared to Thomson parabola data. Electron energy spectra are also collected

¹This work is supported by US DOE/NNSA, performed at LANL, operated by LANS LLC under contract DE-AC52-06NA25396.

James Cobble
Los Alamos National Laboratory

Date submitted: 10 Jul 2014

Electronic form version 1.4