

Abstract Submitted
for the DPP14 Meeting of
The American Physical Society

Reduced model prediction of electron temperature profiles in microtearing-dominated NSTX plasmas¹ S.M. KAYE, W. GUTTENFELDER, R. BELL, S. GERHARDT, B. LEBLANC, R. MAINGI, PPPL, Princeton Univ. — A representative H-mode discharge from the National Spherical Torus Experiment (NSTX) is studied in detail as a basis for a time-evolving prediction of the electron temperature profile using an appropriate reduced transport model. The time evolution of characteristic plasma variables such as β_e , ν_e^* , the MHD α parameter and the gradient scale lengths of T_e , T_i and n_e were examined prior to performing linear gyrokinetic calculations to determine the fastest growing microinstability at various times and locations throughout the discharge. The inferences from the parameter evolutions and the linear stability calculations were consistent. Early in the discharge, when β_e and ν_e^* were relatively low, ballooning parity modes were dominant. As both β_e and ν_e^* increased with time, microtearing became the dominant low- k_θ mode, especially in the outer half of the plasma. There are instances in time and radius where other modes, at higher- k_θ , may be important for driving electron transport. The Rebut-Lallia-Watkins (RLW) electron thermal diffusivity model, which is based on microtearing-induced transport, was used to predict the time-evolving electron temperature across most of the profile. The results indicate that RLW does a good job of predicting T_e for times and locations where microtearing was determined to be important, but not as well when microtearing was predicted to be stable or subdominant.

¹This work has been supported by U.S. Dept of Energy contracts DE-AC02-09CH11466.

S.M. Kaye
PPPL, Princeton Univ.

Date submitted: 10 Jul 2014

Electronic form version 1.4