

Abstract Submitted
for the DPP14 Meeting of
The American Physical Society

Wave-Kinetic Simulations of Lower-Hybrid Turbulence driven by Velocity Ring Instabilities¹ GURU GANGULI, CHRIS CRABTREE, Naval Research Laboratory, LEONID RUDAKOV, Icarus Research Inc., MANISH MITHAIWALA, Naval Research Laboratory — We develop numerical solutions to the wave-kinetic equation in a periodic box including the effects of nonlinear (NL) scattering of Lower-hybrid waves giving the evolution of the wave-spectra in wavenumber space. Simultaneously we solve the particle diffusion equation of both the background plasma particles and the ring ions, due to both linear and nonlinear Landau resonances. At initial times for cold ring ions, an electrostatic beam mode is excited, while the kinetic mode is stable. As the instability progresses the ring ions heat, the beam mode is stabilized, and the kinetic mode destabilizes. When the amplitude of the waves becomes sufficient the lower-hybrid waves are scattered (by either nearly unmagnetized ions or magnetized electrons) into electromagnetic magnetosonic waves [Ganguli et al 2010]. The effect of NL scattering is to limit the amplitude of the waves, slowing down the quasilinear relaxation time and ultimately allowing more energy from the ring to be liberated into waves [Mithaiwala et al. 2011]. The effects of convection out of the instability region are modeled, additionally limiting the amplitude of the waves, allowing further energy to be liberated from the ring [Scales et al., 2012]. Results are compared to recent 3D PIC simulations [Winske and Duaghton 2012].

¹This work is supported by the Naval Research Laboratory base program.

Guru Ganguli
Naval Research Laboratory

Date submitted: 11 Jul 2014

Electronic form version 1.4