

Abstract Submitted
for the DPP14 Meeting of
The American Physical Society

Does a Critical Gradient Exist for Alfvén Eigenmode Induced Fast-Ion Transport?¹ C.S. COLLINS, W.W. HEIDBRINK, U. California Irvine, M.A. VAN ZEELAND, C.C. PETTY, D.C. PACE, General Atomics, B.A. GRIERSON, PPPL — In the critical gradient model, if local energetic particle (EP) drive exceeds the Alfvén eigenmode (AE) stability limit, particles diffuse to flatten the pressure profile until marginal stability is maintained. A key signature is a sudden increase in transport above the critical gradient. In DIII-D, the onset of AE-induced EP transport is examined by modulating the EP pressure profile using an off-axis neutral beam while AE activity gradually diminishes during the current ramp. The time evolution of the EP density profile is measured with fast-ion D_α (FIDA) spectroscopy. During quiescent periods, the FIDA intensity rises and decays approximately linearly during and after the beam pulse, whereas during strong AE activity, the modulated FIDA intensity amplitude and decay rate decrease, suggesting additional AE-induced radial diffusion. Hardware upgrades are underway to increase spatial resolution and accommodate the full D_α spectrum, providing better constraints when comparing to predictive models.

¹Work supported by the US Department of Energy under SC-G903402, DE-FC02-04ER54698, and DE-AC02-09CH11466.

Bill Heidbrink
U. California Irvine

Date submitted: 11 Jul 2014

Electronic form version 1.4