

DPP15-2015-000083

Abstract for an Invited Paper
for the DPP15 Meeting of
the American Physical Society

A new class of three-dimensional ideal-MHD equilibria with current sheets

JOAQUIM LOIZU, Max-Planck-Princeton Center for Plasma Physics

Ideal MHD predicts singular current densities in 3D equilibria with nested flux surfaces: a pressure-driven $1/x$ current density that arises around resonant rational surfaces, and a Dirac δ -function current that develops at those surfaces. Only recently have these currents been computed numerically [1], and we provide details of the calculation. We show that locally-infinite shear (i.e. discontinuous rotational-transform) at the resonant surfaces ensures well-defined solutions. Singularities in the current density are allowed in ideal-MHD, but the current passing through any surface must be finite for any physically-acceptable equilibrium model. While the integral of the δ -current density is finite, the $1/x$ current diverges over certain surfaces. This led to the conclusion that pressure gradients cannot exist in the vicinity of rational surfaces and thus that the possible pressure profiles are either fractal [2] or discontinuous [3]. In this talk, we present a new class of 3D, globally-ideal, MHD equilibria with (i) continuously nested surfaces, (ii) arbitrary, continuous and smooth pressure profiles, (iii) arbitrary, 3D boundaries, (iv) without unphysical currents, and which are (v) analytic functions of the boundary [4]. Examples of such equilibria, computed with the SPEC code, are shown and verified against generalized solutions to Newcomb equation, showing excellent convergence. The results imply that a resonant magnetic perturbation can penetrate all the way into the centre of a tokamak without being shielded at the resonant surface, even within ideal MHD.

[1] J. Loizu, S. Hudson, A. Bhattacharjee and P. Helander, *Phys. Plasmas* 22 022501 (2015)

[2] H. Grad, *Phys. Fluids* 10 (1):137 (1967)

[3] O. Bruno and P. Laurence, *Commun. Pur. Appl. Math.* 49 (7):717-764 (1996)

[4] J. Loizu, S. R. Hudson, A. Bhattacharjee, S. Lazerson and P. Helander, letter submitted to *Phys. Plasmas*