

Abstract Submitted
for the DPP15 Meeting of
The American Physical Society

Heat-Flux Measurements from Collective Thomson-Scattering Spectra R.J. HENCHEN, S.X. HU, J. KATZ, D.H. FROULA, Laboratory for Laser Energetics, U. of Rochester, W. ROZMUS, U. of Alberta — Collective Thomson scattering was used to measure heat flux in coronal plasmas. The relative amplitude of the Thomson-scattered power into the up- and downshifted electron plasma wave features was used to determine the flux of electrons moving along the temperature gradient at three to four times the electron thermal velocity. Simultaneously, the ion-acoustic wave features were measured. Their relative amplitude is used to measure the flux of the return-current electrons. The frequencies of these ion-acoustic and electron plasma wave features provide local measurements of the electron temperature and density. These spectra were obtained at five locations along the temperature gradient in a laser-produced blowoff plasma. These measurements of plasma parameters are used to infer the Spitzer–Härm flux ($q_{\text{SH}} = -\kappa \nabla T_e$) and are in good agreement with the values of the heat flux measured from the scattering-feature asymmetries. This material is based upon work supported by the Department of Energy National Nuclear Security Administration under Award Number DE-NA0001944.

R.J. HENCHEN
Laboratory for Laser Energetics, U. of Rochester

Date submitted: 21 Jul 2015

Electronic form version 1.4