

Abstract Submitted
for the DPP16 Meeting of
The American Physical Society

Current sheet formation in a 3D line-tied plasma¹ YAO ZHOU, YI-MIN HUANG, HONG QIN, AMITAVA BHATTACHARJEE, Princeton University — Recently a variational integrator for ideal MHD in Lagrangian labeling has been developed by discretizing Newcomb’s Lagrangian on a moving mesh using discretized exterior calculus. With the frozen-in equation built-in, the method is free of artificial reconnection, and therefore optimal for studying current sheet formation. Using this method, it is confirmed that the nonlinear solution to the ideal Hahm-Kulsrud-Taylor problem in 2D yields a singular current sheet. We identify it by showing that the equilibrium solution converges with increasing resolution, except where there is singularity. This approach is in contrast to previous studies which use diverging peak current density as sole evidence of current singularity. We then extend the problem to 3D line-tied geometry. The linear solution, which is singular in 2D, is found to be smooth, but pathological when the system is sufficiently long. Accordingly, the nonlinear solution turns out to be smooth for short systems, but tends to become more singular when the system length increases. A resolution to this problem can potentially settle the long-standing controversy over Parker’s conjecture on the formation of current singularity in 3D line-tied geometry.

¹This research was supported by the U.S. DOE under Contract No. DE-AC02-09CH11466.

Yao Zhou
Princeton Plasma Phys Lab

Date submitted: 15 Jul 2016

Electronic form version 1.4