

Abstract Submitted
for the DPP16 Meeting of
The American Physical Society

Investigating the Dynamics of Canonical Flux Tubes¹ JENS VON DER LINDEN, University of Washington, JASON SEARS, Lawrence Livermore National Laboratory, THOMAS INTRATOR², Los Alamos National Laboratory, SETTHIVOINE YOU, University of Washington — Canonical flux tubes are flux tubes of the circulation of a species' canonical momentum. They provide a convenient generalization of magnetic flux tubes to regimes beyond magnetohydrodynamics (MHD). We hypothesize that hierarchies of instabilities which couple disparate scales could transfer magnetic pitch into helical flows and vice versa while conserving the total canonical helicity. This work first explores the possibility of a sausage instability existing on top of a kink as mechanism for coupling scales, then presents the evolution of canonical helicity in a gyrating kinked flux rope. Analytical and numerical stability spaces derived for magnetic flux tubes with core and skin currents indicate that, as a flux tube lengthens and collimates, it may become kink unstable with a sausage instability developing on top of the kink. A new analysis of 3D magnetic field and ion flow data on gyrating kinked magnetic flux ropes from the Reconnection Scaling Experiment tracks the evolution of canonical flux tubes and their helicity. These results and methodology are being developed as part of the Mochi experiment specifically designed to observe the dynamics of canonical flux tubes.

¹This work is supported by DOE Grant DE-SC0010340 and the DOE Office of Science Graduate Student Research Program and prepared in part by LLNL under Contract DE-AC52-07NA27344. LLNL-ABS-697161

²Deceased 3 June 2014.

Jens von der Linden
University of Washington

Date submitted: 15 Jul 2016

Electronic form version 1.4