

Abstract Submitted  
for the DPP17 Meeting of  
The American Physical Society

**Plasma response model to resonant magnetic perturbations**

MARISA ROBERTO, Instituto Tecnológico de Aeronáutica, São José dos Campos, Brazil, ANDRE CARLOS FRAILE JUNIOR, Instituto Tecnológico de Aeronáutica, Instituto de Estudos Avancados, Divisão de Aerodinâmica de Hipersonica, IBERE LUIZ CALDAS, Universidade de São Paulo, Instituto de Física, São Paulo, Brazil — Magnetically confined plasmas in tokamaks usually do not operate under equilibrium conditions and high temperature particles are transported to the reactor wall, causing its erosion. External resonant windings or coils can be used to generate a perturbative magnetic field that modifies particle transport near the plasma border. A pair of helical wires placed at the reactor external wall is used to create a resonant magnetic perturbation (RMP) capable of reducing the amount of particles colliding with the wall. However, the plasma is also affected by RMPs and it modifies the magnetic field lines. In this work, a semi-analytical model in polar toroidal coordinates was developed to study the plasma response, which has been modeled as a helical current sheet at the resonant surface with the condition that the total radial magnetic field vanishes at this surface. This condition is associated to the mitigation of magnetic islands around the resonant surface. Poincaré plots have shown that the plasma response regularizes the magnetic field lines around the resonant surface.

Marisa Roberto  
Instituto Tecnológico de Aeronáutica

Date submitted: 30 Jun 2017

Electronic form version 1.4