

Abstract Submitted
for the DPP17 Meeting of
The American Physical Society

Precursor Events Involving Plasmas Structures

Around Collapsing Black Holes Binaries¹ M. MEDVEDEV, B. COPPI, MIT

— The plasma structures that can exist around black hole binaries can sustain intrinsic plasma collective modes [1] that have characteristic low frequencies related to the particle rotation frequencies around the binary system. As the collapse approaches, with the loss of angular momentum by emission of gravitational waves [2] from the binary system we have suggested [3] that the frequency of the fluctuating component of the gravitational potential can go through that of the intrinsic modes of the surrounding plasma structure and lead to a sharp amplification of them. Then the precursor to the event reported in Ref. [2], tentatively identified by the Agile X- γ -ray observatory [4] may be associated with the high energy radiation emission due to the fields produced by excitation of the proposed plasma modes. M. Tavani is thanked for bringing Ref. [4] to our attention while Ref. [3] was being completed. B. Coppi, Plasma Physics Report, 43, 3, 289297 (2017).

B. Abbot, R. Abbot, T. Abbot et al. Phys. Rev. Lett. 118, 221101 (2017).

B. Coppi and M. Medvedev, MIT-LNS HEP 17/02 June 2016. To be submitted to Physics Letters.

F. Verrecchia, M. Tavani, A. Ursi, et al., to be published in Ap.J. Letters.

¹Sponsored in part by the U.S. DoE.

B. Coppi
MIT

Date submitted: 13 Jul 2017

Electronic form version 1.4