

Abstract Submitted  
for the DPP17 Meeting of  
The American Physical Society

**Development of a Buried Layer Platform at the OMEGA Laser to Study Open L-Shell Spectra from Coronal (non-LTE) Plasmas<sup>1</sup>**  
EDWARD MARLEY, CHARLIE JARROT, MARILYN SCHNEIDER, ELIJAH KEMP, MARK FOORD, ROBERT HEETER, DUANE LIEDAHL, KLAUSE WIDMANN, CHRISTOPHER MAUCHE, GREG BROWN, JAMES EMIG, Lawrence Livermore National Laboratory — A buried layer platform is being developed at the OMEGA laser to study the open L-shell spectra of coronal (non-LTE) plasmas ( $n_e \sim \text{few } 10^{21}/\text{cm}^3$ ,  $T_e \sim 0.8 - 1.2 \text{ keV}$ ) of mid Z materials. Studies have been done using a  $250 \mu\text{m}$  diameter dot composed of a layer of  $1200 \mu\text{m}$  thick Zn between two  $600 \mu\text{m}$  thick layers of Ti, in the center of a  $1000 \mu\text{m}$  diameter,  $13 \mu\text{m}$  thick beryllium tamper. Lasers heat the target from both sides for up to 3 ns. The size of the microdot vs time was measured with x-ray imaging (face-on and side-on). The radiant x-ray power was measured with a low-resolution absolutely calibrated x-ray spectrometer (DANTE). The temperature was measured from the Ti helium-beta complex. The use of this platform for the verification of atomic models is discussed.

<sup>1</sup>This work was performed under the auspices of the U.S. Department of Energy by Lawrence Livermore National Laboratory under Contract DE-AC52-07NA27344.

Edward Marley  
Lawrence Livermore National Laboratory

Date submitted: 13 Jul 2017

Electronic form version 1.4