

Abstract Submitted
for the DPP17 Meeting of
The American Physical Society

Development of a sub-cm high resolution ion Doppler tomography diagnostics for fine structure measurement of guide field reconnection in TS-U¹

HIROSHI TANABE, HIDEYA KOIKE, HIRONORI HATANO, TAKUMI HAYASHI, QINGHONG CAO, SHUNICHI HIMENO, TAISHI KANEDA, MOE AKIMITSU, ASUKA SAWADA, YASUSHI ONO, Graduate School of Frontier Sciences, University of Tokyo — A new type of high-throughput/high-resolution 96CH ion Doppler tomography diagnostics has been developed using "multi-slit" spectroscopy technique for detailed investigation of fine structure formation during high guide field magnetic reconnection. In the last three years, high field merging experiment in MAST pioneered new frontiers of reconnection heating [1]: formation of highly peaked structure around X -point in high guide field condition ($B_t > 0.3T$), outflow dissipation under the influence of better plasma confinement to form high temperature ring structure which aligns with closed flux surface of toroidal plasma, and interaction between ion and electron temperature profile during transport/confinement phase to form triple peak structure ($\tau_{ei}^E \sim 4ms$). To investigate more detailed mechanism with in-situ magnetic measurement, the university of Tokyo starts the upgrade of plasma parameters and spatial resolution of optical diagnostics as in MAST. Now, a new type of high-throughput/high-resolution 96CH ion Doppler tomography diagnostics system construction has been completed and it successfully resolved fine structure of ion heating downstream, aligned with closed flux surface formed by reconnected field. [1] H. Tanabe et al., Phys. Rev. Lett. 115, 215004 (2015)

¹This work was supported by JSPS KAKENHI Grant Numbers 15H05750, 15K14279 and 17H04863.

Hiroshi Tanabe
Graduate School of Frontier Sciences, University of Tokyo

Date submitted: 14 Jul 2017

Electronic form version 1.4