

Abstract Submitted  
for the DPP17 Meeting of  
The American Physical Society

**Electron-Positron Cascade in Magnetospheres of Spinning Black Holes**<sup>1</sup> ALEX L. FORD, KU, BRETT D. KEENAN, LANL, MIKHAIL V. MEDVEDEV, KU MIT — We quantitatively study the stationary, axisymmetric, force-free magnetospheres of spinning (Kerr) black holes (BHs) and the conditions needed for relativistic jets to be powered by the Blandford-Znajek mechanism. These jets could be from active galactic nuclei, blazars, quasars, micro-quasars, radio active galaxies, and other systems that host Kerr BHs. The structure of the magnetosphere determines how the BH energy is extracted, e.g., via Blandford-Znajek mechanism, which converts the BH rotational energy into Poynting flux. The key assumption is the force-free condition, which requires the presence of plasma with the density being above the Goldreich-Julian density. Unlike neutron stars, which in principle can supply electrons from the surface, BH cannot supply plasma at all. The plasma must be generated in situ via an electron-positron cascade, presumably in the gap region. Here we study varying conditions that provide a sufficient amount of plasma for the Blandford-Znajek mechanism to work effectively.

<sup>1</sup>The authors acknowledge DOE partial support via grant DE-SC0016368.

Mikhail Medvedev  
Univ of Kansas

Date submitted: 14 Jul 2017

Electronic form version 1.4