

Abstract Submitted  
for the DPP17 Meeting of  
The American Physical Society

**Laser Induced Fluorescence (LIF) Measurements of Neutral (ArI) and singly-ionized (ArII) Argon in a LargeScale Helicon Plasma<sup>1</sup>**

R. F. KELLY, D. M. FISHER, M. W. HATCH, M. GILMORE, R. H. DWYER, K. MEANY, Univ of New Mexico, Y. ZHANG, University of Washington, T. R. DESJARDINS, Los Alamos National Laboratory — In order to investigate the role of neutral dynamics in helicon discharges in the HelCat (Helicon-Cathode) plasma device at U. New Mexico, a Laser Induced Fluorescence (LIF) system has been developed. The LIF system is based on a >250 mW, tunable diode laser with a tuning range between 680 and 700nm. For neutral Argon, the laser pumps the metastable ( $^2P_{3/2}^0$ )4s level to the ( $^2P_{1/2}^0$ )4p level using 696. 7352 nm light. The fluorescence radiation from decay to the ( $^2P_{1/2}^0$ )4s level at 772. 6333 nm is observed. For singly ionized Argon, the laser pumps the  $3s^23p^4(^3P)3d$  level to the  $3s^23p^4(^3P)4p$  level using 686.3162nm light. The fluorescence radiation from the decay to the  $3s^23p^4(^3P)4s$  level is observed. The system design, and velocity measurements in the axial, azimuthal and radial directions for ArI, and in the axial direction for ArII will be presented.

<sup>1</sup>Supported by U.S. National Science Foundation Award 1500423

R. F. Kelly  
Univ of New Mexico

Date submitted: 14 Jul 2017

Electronic form version 1.4