

Abstract Submitted
for the DPP17 Meeting of
The American Physical Society

Improving Coil Designs for the HSX Stellarator with FOCUS¹

THOMAS KRUGER, Univ of Wisconsin, Madison, CAO XIANG ZHU, Princeton Plasma Physics Laboratory, AARON BADER, LUQUANT SINGH, DAVID ANDERSON, University of Wisconsin, Madison — We use the FOCUS code to generate improved coil sets for the HSX stellarator. FOCUS produces curves in 3D space to best reproduce a target plasma equilibrium. Unlike similar codes, the curves in FOCUS are not constrained to lie on a user-defined 2D surface. Therefore FOCUS can inherently solve problems such as determining the optimum coil-plasma distance for a given equilibrium. By adjusting the relative weights between a) the match to the plasma boundary, and b) the average coil length. We present the results from FOCUS where we attempt to improve the coil set by moving coils further away to reduce coil ripple, decreasing the number of coils to improve accessibility, and better matching the target plasma surface. We also present results of alternative coil designs with helical and saddle coils.

¹Work supported by the US DOE under grant DE-FG02-93ER54222 and UW Sorden account 233PRJ65ZM

Aaron Bader
Univ of Wisconsin, Madison

Date submitted: 14 Jul 2017

Electronic form version 1.4