

Abstract Submitted
for the DPP19 Meeting of
The American Physical Society

Magnetized plasma flow interactions in pulsed-power driven experiments¹ LEE SUTTLE, J.W.D. HALLIDAY, J.D. HARE, D. RUSSELL, E. TUBMAN, V. VALENZUELA-VILLASECA, S.V. LEBEDEV, Imperial College London — The interactions of fast-streaming, magnetized plasmas can result in a wide range of fundamental plasma physics processes such as the formation of MHD shocks, magnetic turbulence, reconnection and wave-particle interactions. We present experiments from a versatile platform, where supersonic plasma flows generated by the ablation of pulsed-power driven wire arrays are used to study a wide range of magnetized plasma interactions [1,2]. The setup allows a control over the global system parameters, including the drive strength, magnetization, magnetic field topology and interaction geometry. The plasma composition (wire material) can also be chosen to vary the collisionality of the plasma and introduce dynamically significant radiative cooling. The detailed structure of the interactions is measured using optical Thomson scattering, multi-color laser interferometry and Faraday rotation diagnostics, providing measurements of the flow velocities, plasma temperature, electron density and magnetic field distributions of the plasma. [1] Suttle et al., PRL 116, 225001 (2016) [2] Burdiak et al., PoP 24, 072713 (2017)

¹Experimental work carried out at the MAGPIE pulsed-power generator, Imperial College London. Supported by DOE Cooperative Agreement DE-NA0003764.

Lee Suttle
Imperial College London

Date submitted: 09 Aug 2019

Electronic form version 1.4