

Abstract Submitted
for the DPP20 Meeting of
The American Physical Society

Hydroscaling and Alpha Heating in High Adiabatic Layered Implosions KEVIN BAKER, OGGIE JONES, STEVE MACLAREN, PATEL PRAV, DEBBIE CALLAHAN, OMAR HURRICANE, DAN CASEY, MATHIAS HOHENBERGER, SHAHAB KHAN, ANNIE KRITCHER, RICCARDO TOMMASINI, EDDIE DEWALD, BRIAN SPEARS, TILO DOEPPNER, ALEX ZYLSTRA, NINO LANDEN, DAN CLARK, CHRIS WEBER, BEN BACHMANN, RICHARD BIONTA, KELLY HAHN, DAVID FITTINGHOFF, DAVID SCHLOSSBERG, Lawrence Livermore Natl Lab, MARIE GATU-JOHNSON, MIT, PETR VOLEGOV, LANL — High adiabatic implosions in inertial confinement fusion (ICF) are designed to be more robust to detrimental plasma and hohlraum physics than their lower adiabatic counterparts. They drive a strong first shock into the ablator as well as into the DT fuel, reducing the sensitivity of the integrated system to uncertainties in shock-timing, preheat, and instabilities in the ablator and at the fuel-ablator interface. The higher adiabatic enables a short pulse which simplifies hohlraum physics by limiting the extent of difficult to model dynamics such as gold bubble expansion, plasma filling of the hohlraum, and stagnation and interpenetration of the wall, capsule and gas interfaces and subsequent laser propagation through those regions. We report on DT layered implosions used to test the level of alpha heating driven in these high adiabatic implosions and the hydroscaling of these implosions between two scales, scaled by $\times 1.125$. We present hydroscalings of the hotspot parameters which are shown to scale differently with the scale factor than no alpha heating analytic theory predicts. *This work was performed under the auspices of the U.S. Department of Energy by Lawrence Livermore National Laboratory under Contract DE-AC52-07NA27344.

Kevin Baker
Lawrence Livermore Natl Lab

Date submitted: 07 Jul 2020

Electronic form version 1.4