

Abstract Submitted
for the GEC09 Meeting of
The American Physical Society

Electron Density Measurements in a Supersonic Flowing Ar/H₂/Air Discharge M. NIKOLIC, D.J. DRAKE, P. LAURIN, S. POPOVIC, L. VUSKOVIC, Department of Physics, Old Dominion University — Local measurements of the electron density in high pressure discharges have commonly been performed through use of an electrical probe, such as a Langmuir probe, or by the Stark broadening of the hydrogen lines. However in supersonic flowing discharges electrical probes can cause shocks to form, which is unwanted. In addition, these types of discharges do not often contain the hydrogen needed to determine the electron density through Stark broadening where the measurements are hampered by the lack of intensity and breadth of the hydrogen Balmer lines. An alternative approach is to use the intensity of the rotational bands of the N₂ second positive system. We performed detailed measurements of the population densities of the N₂ C³Π_u-B³Π_g system and the hydrogen Balmer lines in a supersonic flow of weakly ionized Ar/H₂/Air. Gases were premixed in the stagnation chamber at room temperature by adding up to 10% hydrogen and up to 45% air to pure argon. A cylindrical cavity was used to sustain a discharge in the pressure range of 100-700 Pa. Absolute emission spectroscopy was used to determine the gas temperature in the flow from the N₂ system. Comparison was made between the results obtained from the N₂ band intensity technique and Stark broadening of the hydrogen Balmer lines.

M. Nikolic
Department of Physics, Old Dominion University

Date submitted: 11 Jun 2009

Electronic form version 1.4