

Abstract Submitted
for the GEC10 Meeting of
The American Physical Society

Electron attachment to CF_2 and preliminary results for C_2F_5 ¹

THOMAS FIELD, Queens University Belfast, KAROLA GRAUPNER, Loughborough University, CHRISTOPHER MAYHEW, Birmingham University, UK, SEAN HAUGHEY, Queens University Belfast, JUDITH LANGER, TU Berlin, EUGEN ILLENBERGER, Frei University Berlin — Experiments have been performed to investigate electron attachment to highly reactive molecular fragments CF_2 and C_2F_5 . CF_2 was generated in a plasma discharge of C_3F_6 (perfluoropropene) and by reaction of hydrogen atoms with CF_3I . The C_2F_5 radical was generated by reaction of $\text{C}_2\text{F}_5\text{I}$ with hydrogen atoms. Electron attachment experiments were made with ERIC, which includes a low energy pulsed trochoidal electron gun and a small linear time-of-flight mass spectrometer (Field et al., J. Phys. B 38 (2005) 255). In the case of CF_2 no electron attachment was observed, despite theoretical prediction of a significant dissociative electron attachment cross section (Rozum et al., J. Phys. Chem. Ref. Data 35 (2006) 267). An upper limit for the dissociative electron attachment cross section of CF_2 has been estimated as $5 \times 10^{-4} \text{ \AA}^2$ at 1.8 eV, which is the thermodynamic threshold for F^- formation from CF_2 . In the case of C_2F_5 the data are under analysis and results will be presented at the meeting.

¹The authors gratefully acknowledge the support of EPSRC, the UK Non-Thermal Plasma Network and the european ITS-LEIF network.

Thomas Field
Queens University Belfast

Date submitted: 14 Jun 2010

Electronic form version 1.4