

Abstract Submitted
for the GEC12 Meeting of
The American Physical Society

Simulation of a capacitively coupled oxygen discharge using the oopd1 particle-in-cell Monte Carlo code J.T. GUDMUNDSSON, Shanghai Jiao Tong University, M.A. LIEBERMAN, University of California at Berkeley — The oopd1 particle-in-cell Monte Carlo (PIC-MC) code is used to simulate a capacitively coupled discharge in oxygen. oopd1 is a one-dimensional object-oriented PIC-MC code in which the model system has one spatial dimension and three velocity components. It contains models for planar, cylindrical, and spherical geometries and replaces the XPDx1 series, which is not object-oriented. The revised oxygen model includes, in addition to electrons, the oxygen molecule in ground state, the oxygen atom in ground state, the negative ion O^- , and the positive ions O^+ and O_2^+ . The cross sections for the collisions among the oxygen species have been significantly revised from earlier work using the XPDP1 code. Here we explore the electron energy distribution function (EEDF), the ion energy distribution function (IEDF) and the density profiles for various pressures and driving frequencies. In particular we investigate the IEDF for both O^+ and O_2^+ ions for various discharge conditions. We explore the role of fragmentation due to collisions of high energy O_2^+ ions with oxygen molecules its influence on the IEDF for O^+ -ions.

Jon Gudmundsson
Shanghai Jiao Tong University

Date submitted: 26 Jun 2012

Electronic form version 1.4