

Abstract Submitted  
for the GEC16 Meeting of  
The American Physical Society

**Non-locality, adiabaticity, thermodynamics and electron energy probability functions.** RODERICK BOSWELL, YUNCHAO ZHANG, CHRISTINE CHARLES, Australian Natl Univ, KAZUNORI TAKAHASHI, Tohoku University — Thermodynamic properties are revisited for electrons that are governed by nonlocal electron energy probability functions in a plasma of low collisionality. Measurements in a laboratory helicon double layer experiment have shown that the effective electron temperature and density show a polytropic correlation with an index of  $\gamma_e = 1.17 \pm 0.02$  along the divergent magnetic field, implying a nearly isothermal plasma ( $\gamma_e = 1$ ) with heat being brought into the system. However, the evolution of electrons along the divergent magnetic field is essentially an adiabatic process, which should have a  $\gamma_e = 5/3$ . The reason for this apparent contradiction is that the nearly collisionless plasma is very far from local thermodynamic equilibrium and the electrons behave nonlocally. The corresponding effective electron enthalpy has a conservation relation with the potential energy, which verifies that there is no heat transferred into the system during the electron evolution. The electrons are shown in nonlocal momentum equilibrium under the electric field and the gradient of the effective electron pressure. The convective momentum of ions, which can be assumed as a cold species, is determined by the effective electron pressure and the effective electron enthalpy is shown to be the source for ion acceleration. For these nearly collisionless plasmas, the use of traditional thermodynamic concepts can lead to very erroneous conclusions regarding the thermal conductivity.

Roderick Boswell  
Australian Natl Univ

Date submitted: 09 Jun 2016

Electronic form version 1.4