

Abstract Submitted
for the GEC19 Meeting of
The American Physical Society

Stopping of proton beam in hydrogen¹ ALISHER KADYROV, JACKSON BAILEY, ILKHOM ABDURAKHMANOV, IGOR BRAY, Curtin University, AKRAM MUKHAMEDZHANOV, Texas AM University — The stopping cross section for protons passing through hydrogen is calculated for the energy range between 10 keV and 3 MeV. Both the positive and neutral charge states of the projectile are accounted for. The two-center convergent close-coupling method is used to model proton collisions with hydrogen. In this approach, electron-capture channels are explicitly included by expanding the scattering wave function in a basis of both target and projectile pseudostates. Hydrogen collisions with hydrogen are modeled using two methods: the single-center convergent close-coupling approach is used for the calculation of one-electron processes, while two-electron processes are calculated using the Born approximation. The aforementioned approaches are also applied to the calculation of the charge-state fractions. These are then used to combine the proton-hydrogen and hydrogen-hydrogen stopping cross sections to yield the total stopping cross section for protons passing through hydrogen.

¹This work was supported by the Australian Research Council. A.S.K. acknowledges partial support from U.S. NSF under Award No. PHY-1415656. A.M.M. was supported by U.S. DOE Grant No. DEFG02-93ER40773, NNSA Grant No. DE-NA0003841, and U.S. NSF Grant No. PHY-1415656.

Alisher Kadyrov
Curtin University

Date submitted: 29 May 2019

Electronic form version 1.4