

Abstract Submitted
for the HAW05 Meeting of
The American Physical Society

Study of J/Ψ at high temperatures in anisotropic lattice QCD
HIDEAKI IIDA, NORIYOSHI ISHII, Tokyo Institute of Technology, HIDEO SUG-
ANUMA, Kyoto University — The high-temperature J/Ψ mode above the QCD
critical temperature T_c is studied using anisotropic quenched lattice QCD. We aim
to clarify whether the J/Ψ state above T_c is a localized resonance state or a $c\bar{c}$
scattering state. We investigate $c\bar{c}$ modes for $1.1T_c < T < 2.1T_c$ using the $O(a)$ -
improved Wilson quark action at $\beta = 6.10$ with renormalized anisotropy $a_s/a_t = 4$.
To distinguish localized states and scattering states, we calculate the $c\bar{c}$ correlators
on finite lattices with different spatial boundary conditions, i.e., the periodic and
the anti-periodic boundary condition. (Note that the $c\bar{c}$ threshold is raised up in
the anti-periodic boundary condition.) As a result, almost no energy difference is
found for the lowest J/Ψ mode between the periodic and the anti-periodic boundary
conditions at for $1.1T_c < T < 2.1T_c$. This fact indicates that the lowest J/Ψ mode
can survive as a localized state even above T_c in quenched QCD.

Hideaki Iida
Tokyo Institute of Technology

Date submitted: 24 May 2005

Electronic form version 1.4