

Abstract Submitted  
for the HAW05 Meeting of  
The American Physical Society

**Chiral Three-Nucleon Interaction and Structure of Light Nuclei<sup>1</sup>**

V.G. GUEORGUIEV, PETR NAVRATIL, ERICH ORMAND, Lawrence Livermore National Laboratory — Modern high-precision nucleon-nucleon (NN) interactions derived from the two nucleon system fail to describe systems of more than two-nucleons; therefore, they need to be augmented by three-nucleon interactions (TNI). Compared to the NN-interaction, the form and parameters of the TNI are much less explored. Chiral perturbation theory ( $\chi PT$ ) at next-to-next-to-leading order (N<sup>2</sup>LO) predicts three types of TNI terms. A two-pion exchange term, which is frequently used, has parameters determined from the two-nucleon  $\chi PT$ -potential and/ or pion-nucleon scattering data. The other two terms, the one-pion exchange with NN-contact term and the NNN-contact term, are accompanied by undetermined low-energy constants. We include all three TNI terms in *ab initio* no-core shell model (NCSM) calculations. The two undetermined TNI constants are fit to binding energies of *s*-shell nuclei. The impact of the N<sup>2</sup>LO TNI on the nuclear structure is then studied in NCSM calculations for *p*-shell nuclei, such as <sup>10,11</sup>B that are particularly known to be poorly described by the modern NN potentials.

<sup>1</sup>This work was performed under the auspices of the U. S. Department of Energy by the University of California, Lawrence Livermore National Laboratory under contract No. W-7405-Eng-48. Support from DOE grant SCW0498 is acknowledged.

V.G. Gueorguiev  
Lawrence Livermore National Laboratory

Date submitted: 25 May 2005

Electronic form version 1.4