

Abstract Submitted  
for the HAW05 Meeting of  
The American Physical Society

**Laser Spectroscopic Determination of the Nuclear Charge Radius of  ${}^6\text{He}$  and  ${}^8\text{He}$** <sup>1</sup> P. MUELLER, L.-B. WANG, K. BAILEY, J.P. GREENE, D. HENDERSON, R.J. HOLT, R.V.F. JANSSENS, C.L. JIANG, Z.-T. LU, T.P. O'CONNOR, R.C. PARDO, K.E. REHM, J.P. SCHIFFER, X.D. TANG, Argonne National Laboratory, G.W.F. DRAKE, University of Windsor — Laser spectroscopic measurements of atomic isotope shifts provide unique access to the nuclear charge distribution of short-lived isotopes. The isotopes of interest for this study are  ${}^6\text{He}$  ( $t_{1/2} = 807$  ms) and  ${}^8\text{He}$  ( $t_{1/2} = 119$  ms), which exhibit a loosely bound neutron halo around an  $\alpha$ -like core. Charge radii measurements of both isotopes provide corroboration for their halo structure and test nuclear structure theories of light nuclei. We have performed high-resolution laser spectroscopy on individual radioactive helium atoms captured in a magneto-optical trap. This technique enabled us to accurately measure the isotope shift between  ${}^6\text{He}$  and  ${}^4\text{He}$  in a selected atomic transition. Based on this result and precision atomic theory calculation of helium, the root-mean-square charge radius of the  ${}^6\text{He}$  nucleus could be determined to be 2.054(14) fm [1]. Currently, we are working to expand this technique to also measure the nuclear charge radius of  ${}^8\text{He}$ , and we will report on first results from this experiment. [1] L.-B. Wang *et al.*, PRL 93, 142501 (2004)

<sup>1</sup>This work was supported by the U.S. Department of Energy, Office of Nuclear Physics, under Contract No. W-31-109-ENG-38

Peter Mueller  
Argonne National Laboratory

Date submitted: 25 May 2005

Electronic form version 1.4