

Abstract Submitted
for the HAW05 Meeting of
The American Physical Society

Nucleon Motion Effects for Hyperon Production Reactions

SHINTARO HASHIMOTO, Department of Physics, Kyushu University, MICHIO KOHNO, Physics Division, Kyushu Dental College, KAZUYUKI OGATA, Department of Physics, Kyushu University, YUKINOBU WATANABE, Department of Advanced Energy Engineering Science, Kyushu University, MITSUJI KAWAI, Department of Physics, Kyushu University — Single-particle potentials of hyperon, e.g. Λ , Σ , Ξ , in nuclei are determined from analysis of $^{32}\text{Si}(\pi^\pm, K^+)$ at 1.2 GeV/c and $^{12}\text{C}(K^-, K^+)$ at 1.65 GeV/c. We use the semiclassical distorted wave (SCDW) model to accurately describe these reactions. As an important advantage of the use of the SCDW model, the Fermi motion of a struck nucleon in the target nucleus is explicitly taken into account. Consequently, the Σ - ^{32}Si potential derived from $^{32}\text{Si}(\pi^-, K^+)$ with the SCDW model is significantly different from a result of previous analysis, where a rather simple reaction-model was adopted. In the present paper we discuss effects of the Fermi motion of a struck target-nucleon on the energy spectrum of the outgoing particle in detail and aim to clarify the reaction mechanisms of the inclusive (π^\pm, K^+) and (K^-, K^+) reactions. We will also discuss the role of multistep processes, which can easily be included in the SCDW model analysis, in the study of the hyperon single-particle potential.

Shintaro Hashimoto
Department of Physics, Kyushu University

Date submitted: 03 Jun 2005

Electronic form version 1.4