

Abstract Submitted
for the HAW05 Meeting of
The American Physical Society

Selective Population Shown by Five-Particle Transfer Reactions¹

A.M. CRISP, N. KEELEY, K.W. KEMPER, O. MOMOTYUK, B.T. ROEDER, M. WIEDEKING, Florida State University, J. LIENDO, Simon Bolivar University, F. MARÉCHAL, Strasbourg, France, K. RUSEK, Warsaw, Poland — Spectra and angular distributions were obtained for the reactions $^{16}\text{O}(\text{d,p})^{17}\text{O}$, $^{14}\text{N}(^7\text{Li},\alpha)^{17}\text{O}$, $^{14}\text{N}(^6\text{Li},^3\text{He})^{17}\text{O}$, $^{12}\text{C}(^7\text{Li},\text{d})^{17}\text{O}$, and $^{12}\text{C}(^6\text{Li},\text{p})^{17}\text{O}$. Excitation functions were also measured for $^{12}\text{C}(^7\text{Li},\text{d})$ in the energy range 32-35 MeV and for $^{12}\text{C}(^6\text{Li},\text{p})$ in the energy range 26-32 MeV. These data were collected using the FSU Tandem/LINAC accelerator. Selective population of states in ^{17}O was observed in each of these reactions, and the states populated in each reaction type did not vary with energy. This result indicates a direct, single-step transfer in each case. Further evidence for single-step transfer in the five-particle transfer reactions is provided by DWBA calculations performed on the selectively populated levels. In addition, characterization of particular ^{17}O levels above 11 MeV will be presented. The most strongly excited state is at 11.82 MeV and its spin has been determined to be 7/2.

¹Work supported by the National Science Foundation and the State of Florida

A.M. Crisp
Florida State University

Date submitted: 28 Jun 2005

Electronic form version 1.4