

Abstract Submitted
for the HAW14 Meeting of
The American Physical Society

Coupling Algorithm for $\text{Sp}(3, \mathbb{R})$ Irreducible Representations¹

JAMES ST GERMAINE-FULLER, Grinnell College, ANNA MCCOY, MARK CAPRIO, Notre Dame University — The nuclear symplectic model based on $\text{Sp}(3, \mathbb{R})$ – the smallest algebra that contains both the shell model Hamiltonian and the rotor algebra – connects the microscopic shell model to collective rotational behavior and naturally extends the Elliot $\text{SU}(3)$ model to multiple shells. However, $\text{Sp}(3, \mathbb{R})$ is only an approximate symmetry of the nucleus, where the symmetry can be broken by spin-orbit interactions, tensor force interactions, and pairing. The Hamiltonians in most physical situations will break $\text{Sp}(3, \mathbb{R})$ symmetry, causing their eigenstates to become linear combinations of symplectic irreducible representations (irreps). Calculations with those eigenstates will then involve multiple irreps. We report a computer algorithm for coupling symplectic irreps that should assist in performing such multi-irrep calculations and facilitate computing symplectic coupling coefficients.

¹Supported by the US NSF under grant NSF-PHY05-52843, the US DOE under grant DE-FG02-95ER-40934, and the Research Corporation for Science Advancement under a Cottrell Scholar Award.

James St Germaine-Fuller
Grinnell College

Date submitted: 24 Jul 2014

Electronic form version 1.4