

Abstract Submitted
for the MAR05 Meeting of
The American Physical Society

Probing of Charging and Reactivity of Supported Gold Nanoclusters: Bonding of CO to Gold Octamers Adsorbed on Magnesia
BOKWON YOON, HANNU HÄKKINEN, UZI LANDMAN, School of Physics, Georgia Institute of Technology, Atlanta, GA 30332-0430, A. WÖRZ, J.-M. ANTONIETTI, S. ABBET, K. JUDAI, U. HEIZ, Departement Chemie, Lehrstuhl für Physikalische Chemie I, Technische Universität München, 85747 Garching, Germany — Gold octamers supported on Mg(001) are the smallest clusters to catalyze the low-temperature oxidation of CO to carbon dioxide, while clusters deposited on close-to-perfect magnesia surfaces remain chemically inert. Charging of the supported clusters plays a key role in promoting their chemical activity. Direct evidence for enhanced electron transfer from the F-centers to the anchored clusters is provided here through infra-red measurements, under reaction conditions, of the stretch vibration of CO adsorbed on mass-selected gold octamers soft-landed on close-to-perfect or defect-rich MgO(001). The experiments agree with quantum ab-initio calculations that predict a red-shift of the CO vibration when the molecules are adsorbed on gold octamers anchored to F-centers. This results from electron back-donation to the CO anti-bonding orbital that weakens the intra-molecular bond.

Bokwon Yoon
School of Physics, Georgia Institute of Technology, Atlanta, GA 30332-0430

Date submitted: 22 Dec 2004

Electronic form version 1.4