

Abstract Submitted  
for the MAR05 Meeting of  
The American Physical Society

**Current Percolation Characteristics of IBAD Coated Conductors**

G. YOU, K.R. BARRACA, L.B. WANG, C. KWON, Department of Physics and Astronomy, California State University Long Beach — We have investigated IBAD coated conductors by variable temperature scanning laser microscopy (VTSLM) to map how current flows in superconducting transition and superconducting region. We examined various factors that make non-uniform current distribution and found that the current flow was affected by the grain boundary network, the physical bending of samples, and the surface contaminations. On the basis of data analysis, we deduced  $65\mu\text{m} \sim 150\mu\text{m}$  as percolation cluster feature sizes. Below the critical temperature, we have investigated  $J_c$  to establish the relationship of the  $T_c$  and  $\delta V$  and  $J_c$ , and we conclude that lower  $T_c$  and higher  $\delta V$  area has lower  $J_c$ . We also found several circular shapes of about  $100\mu\text{m} \times 130\mu\text{m}$  diameter from a sample. Those dots had definitely higher  $T_c$  than the surrounding area and created non-uniform current flow. In this presentation, we will report more about VTSLM images around the dots and the characteristics of current bottleneck features which are related with the lower  $J_c$ .

Chuhe Kwon  
Department of Physics and Astronomy  
California State University Long Beach

Date submitted: 03 Dec 2004

Electronic form version 1.4