MAR06-2005-001470

Abstract for an Invited Paper for the MAR06 Meeting of the American Physical Society

Structure Determination of Noble Metal Clusters by Trapped Ion Electron Diffraction

DETLEF SCHOOSS, Institute for Nanotechnology, Forschungszentrum Karlsruhe

The structures of noble metal cluster ions have been studied by the recently developed technique of trapped ion electron diffraction $(TIED)^1$. In brief, cluster ions are generated by a magnetron sputter source and injected into a cooled (95 K) quadrupole ion trap. After mass selection and thermalization, the trapped ions are irradiated with a 40 keV electron beam. The resulting diffraction pattern is integrated with a CCD detector. The assignment of the structural motif is done via a comparison of the experimental and simulated scattering function, calculated from density functional theory structure calculations.

The structures of mass selected silver cluster cations Ag_{19}^+ , Ag_{38}^+ , Ag_{55}^+ , Ag_{59}^+ , Ag_{75}^+ and Ag_{79}^+ have been investigated². The resulting experimental data are best described by structures based on the icosahedral motif, while closed packed structures could be ruled out. Additionally, we present a comparison of the structures of $Cu_{20}^{+/-}$, $Ag_{20}^{+/-}$ and $Au_{20}^{+/-}$. Our findings show unambiguously that the structure of Au_{20}^- is predominantly given by a tetrahedron in agreement with the results of L.S. Wang et al.³ In contrast, structures of Ag_{20}^- and Cu_{20}^- based on the icosahedral motif agree best with the experimental data. Small structural differences between the charge states are observed. The possibilities and limitations of the TIED method are discussed.

(1) M. Maier-Borst, D. B. Cameron, M. Rokni, and J. H. Parks, Physical Review A 59 (5), R3162 (1999); S. Krückeberg,

D. Schooss, M. Maier-Borst, and J. H. Parks, Physical Review Letters 85 (21), 4494 (2000).

(2) D. Schooss, M.N. Blom, B. v. Issendorff, J. H. Parks, and M.M. Kappes, Nano Letters 5 (10), 1972 (2005).

(3) J. Li, X. Li, H. J. Zhai, and L. S. Wang, Science 299, 864 (2003)