

Abstract for an Invited Paper
for the MAR06 Meeting of
The American Physical Society

What is the “glue” for high temperature superconductivity?

TONICA VALLA, Brookhaven National Laboratory

It has been 20 years since the high-temperature superconductivity (HTSC) was discovered in $\text{La}_{2-x}\text{Ba}_x\text{CuO}_4$, by Bednorz and Müller. Since then, many different HTSC compounds, all containing copper-oxide planes, were synthesized and HTSC became one of the most studied problems in science. However, the mechanism of HTSC still remains unknown. Recent advancements in Angle-Resolved Photoemission Spectroscopy (ARPES) have enabled a direct probing of effects of interactions between electrons and different bosonic excitations in a system, raising the expectations that a “pairing boson” responsible for HTSC could finally be identified. Here, we will present the study of single-particle excitations in two different cuprates: $\text{Bi}_2\text{Sr}_2\text{CaCu}_2\text{O}_{8+\delta}$, a high-temperature superconductor with $T_C = 91$ K and $\text{La}_{1.875}\text{Ba}_{0.125}\text{CuO}_4$, a system with suppressed superconductivity ($T_C \leq 2.5$ K) due to the spin/charge ordering. The extracted self-energies and single-particle gaps will be compared and possible coupling mechanisms will be discussed. This work was supported by the DOE under contract number DE- AC02-98CH10886.