

Abstract Submitted  
for the MAR06 Meeting of  
The American Physical Society

**Resolving the Laughlin Paradox** TALBOT CHUBB, Physicist Consultant, 5023 N. 38th St., Arlington, VA 22207 — For paired Bloch electrons in a metal not subject to Pauli exclusion, the 2-electron Hamiltonian has the form

$$H = \frac{-\hbar^2}{4m_e} \Delta_{cm} + (2e)U_{lattice}(r_{cm}, N_{cell}) + \frac{e^2}{(N_{cell}r_{12})} - \frac{\hbar^2}{3m_e} \Delta_{12},$$

where  $r_{cm} = r_1 + r_2$ ,  $r_{12} = r_1 - r_2$ , and  $r_1$  and  $r_2$  are position vectors in configuration space, involving independent Bravais vectors  $R_1$  and  $R_2$ , such that  $R_1 - R_2 = R_{12}$  is an independent Bravais lattice vector, and  $N_{cell}$  is the number of mutually shared potential wells over which the 2 electrons are coherently partitioned with entangled local density maxima. At large  $N_{cell}$ , the magnitude of term 3  $\ll$  the magnitude of term 1. When coordinate exchange symmetry is satisfied and energy minimized, term 3 cancels term 1 at  $r_{12} = 0$ , eliminating the singularity in the wave equation, thereby resolving Laughlin's paradox<sup>1</sup>

<sup>1</sup>R.B. Laughlin, "A Different Universe", (Basic Books, Cambridge MA, 2005) pp. 84-85.

Scott Chubb  
Naval Research Laboratory

Date submitted: 05 Dec 2005

Electronic form version 1.4