

Abstract Submitted  
for the MAR06 Meeting of  
The American Physical Society

**Quantum state control of atoms in microscopic optical traps<sup>1</sup>**

MARK SAFFMAN, DENIZ YAVUZ, MARIE DELANEY, University of Wisconsin, PASAD KULATUNGA, Hobart and William Smith Colleges, TODD JOHNSON, ERICH URBAN, THOMAS HENAGE, NICHOLAS PROITE, THAD WALKER, University of Wisconsin — We present recent progress in loading and manipulation of neutral atoms in microscopic optical traps. Single Rb atoms are loaded into far off resonant optical traps from a background vapor of cold atoms. Tightly focused optical beams are used to perform two-photon stimulated Raman rotations between hyperfine qubit states. We demonstrate qubit rotations at a rate of 1.4 MHz, 1 ms coherence time, and individual site addressing with crosstalk at the level of  $10^{-3}$ . These results are a significant step towards quantum computing using optically trapped neutral atoms. We discuss work in progress aimed at observing strong, angle independent dipole-dipole interactions for fast two-qubit gates using microwave dressing of Rydberg states. We demonstrate two-photon coherent excitation of Rydberg levels by a  $5s_{1/2} - 5p_{3/2} - nd_{5/2}$  sequence. The possibility of dipole-dipole interactions without angular zeroes will be important for gates, as well as for coupling to mesoscopic qubits to enable transmission of quantum states.

<sup>1</sup>This work is funded by the NSF and the Army Research Office.

Mark Saffman  
University of Wisconsin

Date submitted: 02 Dec 2005

Electronic form version 1.4