

Abstract Submitted
for the MAR06 Meeting of
The American Physical Society

Polaron transport in triphenylene-based discotic liquid crystals

VOLODIMYR DUZHKO, Department of Physics, Case Western Reserve University, OH 44106, ALEXANDER N. SEMYONOV, ROBERT J. TWIEG, Department of Chemistry, Kent State University, OH 44242, KENNETH D. SINGER, Department of Physics, Case Western Reserve University, OH 44106 — We report on the investigation of charge carrier transport in the columnar phase of the triphenylene-based discotic liquid crystal, hexapentyloxytriphenylene, by the time-of-flight technique. The hole mobility was found to be temperature and electric field dependent with a maximum value of $2 \times 10^{-3} \text{ cm}^2/\text{Vs}$. Its temperature dependence is described by a T^{-n} power law, with an electric field dependent n varying from 2.5 to 4.5 corresponding to electric field values from $5 \times 10^4 \text{ V/cm}$ to $5 \times 10^3 \text{ V/cm}$, respectively. The drift velocity of charge carriers is a linear function of the electric field for small fields below $5 \times 10^4 \text{ V/cm}$ and tends to saturation at strong fields. These results are interpreted in the framework of correlated polaron motion as described by the non-adiabatic low-temperature limit of Holstein's small polaron theory. The applicability of the Holstein polaron model to triphenylene-based materials showing thermally activated or temperature-independent carrier mobilities will be discussed.

Volodymyr Duzhko
Department of Physics, Case Western Reserve University, OH 44106

Date submitted: 16 Jan 2006

Electronic form version 1.4