

Abstract Submitted
for the MAR06 Meeting of
The American Physical Society

Anomalous Slow Relaxation of the Magnetization in $Y_{1-x}Pr_xBa_2Cu_3O_{7-\delta}$ ¹ P. GYAWALI, V. SANDU², C. C. ALMASAN, Kent State University, B. J. TAYLOR, M. B. MAPLE, University of California at San Diego — We have investigated the time t evolution of the irreversible magnetization M_{irr} in a series of single crystals of $Y_{1-x}Pr_xBa_2Cu_3O_{7-\delta}$, $x = 0.13$ ($T_c = 82$ K), $x = 0.34$ ($T_c = 50$ K), and $x = 0.47$ ($T_c = 34$ K), all displaying a second peak in magnetization. In all cases, $M_{irr}(t)$ follows the well known law of relaxation, $M_{irr}(t) \propto [\mu k_B T / U_0 \ln(t/t_0)]^{-1/\mu}$. For fields/temperatures lower than the corresponding values of the second magnetization peak, μ is anomalously large, $2.5 \leq \mu \leq 4$, in contrast with theoretical predictions which gives a maximum value of 2.5, and with data reported for other cuprates. These large μ values in the vortex glass state, which give rise to a low relaxation rate, occur for all Pr doping and could be due to the presence of Pr ions. In contrast, at fields/temperatures above the second magnetization peak, μ decreases to values expected from theories of plastic vortex liquid and validated by most experimental reports.

¹This research was supported by the NSF under Grant No DMR 0406471 at KSU and the USDOE under Grant No DE-FG03-86ER-45230 at UCSD

²Permanent address: National Institute of Materials Physics, Bucharest-Magurele, Romania

Parshu Gyawali
Kent State University

Date submitted: 30 Nov 2005

Electronic form version 1.4