Abstract Submitted
for the MAR06 Meeting of
The American Physical Society

Observation of the pseudogap features in overdoped Bi2Sr2CaCu2O8+d by breakjunction tunneling spectroscopy[1] Y. XUAN, H. J. TAO, Z. Z. LI, B. R. ZHAO, Z. X. ZHAO, National Laboratory for Superconductivity, Institute of Physics and Center for Condensed Mater Physics, Chinese Academy of Sciences, China, C.T. LIN, Max-Planck-Institut für Festkörperforschung, 70569 Stuttgart, Germany — Tunneling measurements of Bi2Sr2CaCu2O8+d single crystals with different oxygen doping have been carried out by using break-junction technique to study the pseudogap in the overdoped regime. The normal-state pseudogap has been observed clearly in the three overdoped crystals with Tc = 90 K, 82 K, and 72 K, respectively. Furthermore, direct measurements on the pseudogap opening temperature T*, by tracing the disappearance of the tunneling conductance peak, reveal that the relationship of T* versus hole concentration is linear and its extrapolation just crosses the end of the superconducting phase boundary.

[1]This work was Supported by a grant for State Key Program for Basic Research of China, and the National Center for Research and Development on Superconductivity.

National Laboratory for Superconductivity, Institute of Physics and Center for Condensed Mater Physics, Chin

Date submitted: 06 Dec 2005 Electronic form version 1.4