

Abstract Submitted  
for the MAR06 Meeting of  
The American Physical Society

**Determination of  $t_{2g}$  Complex in  $\text{Na}_x\text{CoO}_2$  by Angle-Resolved Photoemission Spectroscopy** HONGBO YANG, Brookhaven National Laboratory, JIHUA MA, ZHIHUI PAN, Boston College, ALEXEI FEDOROV, Lawrence Berkeley National Laboratory, JONGYIN JIN, BRIAN SALES, DAVID MANDRUS, Oak Ridge National Laboratory, ZIQIANG WANG, HONG DING, Boston College — A systematic investigation on  $t_{2g}$  complex in  $\text{Na}_x\text{CoO}_2$  is carried by ARPES. By taking advantage of the matrix element effect, we recover and trace the dispersion of all three  $t_{2g}$  bands for different Na concentration ( $x = 0.3 \sim 0.7$ ). The dispersion “kink” we observed before is found to be caused by the hybridization between  $a_{1g}$  and  $e'_g$  bands. This band hybridization forms a narrow band near the Fermi level, which causes a large mass renormalization. The width of the non-hybridized bands is about half of the one predicted by LDA calculations, which is possibly due to the correlation effect.

Hongbo Yang  
Brookhaven National Laboratory

Date submitted: 04 Jan 2006

Electronic form version 1.4