

Abstract Submitted
for the MAR07 Meeting of
The American Physical Society

The pseudogap phase in $(\text{TaSe}_4)_2\text{I}$ ANDRAS VANYOLOS, BALAZS DORA, ATTILA VIROSZTEK, Department of Physics, Budapest University of Technology and Economics, 1521 Budapest, Hungary — We have developed the mean-field theory of coexisting charge-density wave (CDW) and unconventional charge-density wave (UCDW). The double phase transition manifests itself in the thermodynamic quantities and in the magnetic response, such as spin susceptibility and nuclear spin-lattice relaxation rate. Our theory qualitatively applies to the quasi-one dimensional CDW material $(\text{TaSe}_4)_2\text{I}$. This material exhibits peculiar properties: above the CDW transition temperature T_c , thermal fluctuations were found to die out rapidly, but robust pseudogap behavior is still detected. Namely, the experimental findings include: (i) sharp increase of the static spin susceptibility above T_c , (ii) smooth increase of the spin-lattice relaxation rate above T_c , (iii) as opposed to conventional CDW, no sharp feature in the spin-lattice relaxation rate below T_c . We have found that our coexisting CDW+UCDW model qualitatively describes these observed properties of $(\text{TaSe}_4)_2\text{I}$. Direct calculations for the magnetic response are shown to evidence the agreement. We also argue, that the fluctuations around T_c are suppressed due to the presence of the “hidden” UCDW phase, which partially gaps the Fermi surface, and causes non-Fermi-liquid (pseudogap) behavior.

Andras Vanyolos
Department of Physics, Budapest University of Technology
and Economics, 1521 Budapest, Hungary

Date submitted: 07 Nov 2006

Electronic form version 1.4