

Abstract Submitted
for the MAR07 Meeting of
The American Physical Society

Magnetism **of**
Rubidium Cobalt Hexacyanoferrate Nanoparticles.¹ D.M. PAJEROWSKI,
M.W. MEISEL, Department of Physics, University of Florida, F.A. FRYE, D.R.
TALHAM, Department of Chemistry, University of Florida — Although photoinduced magnetism in nanoparticles of Prussian blue analogs has been reported, these samples are superparamagnetic. We have generated and characterized nanoparticles of $\text{Rb}_j\text{Co}_k[\text{Fe}(\text{CN})_6]_l \cdot n\text{H}_2\text{O}$, which exhibit photoinduced magnetism and, for the largest particles, long-range ferrimagnetism with finite coercive fields. The synthesis involves the variation of the concentration of the poly(vinylpyrrolidone), PVP, the encapsulating polymer, which controls the resulting particle sizes. From HR-TEM, the particle size distributions have been obtained for four batches of samples, with mean diameters ranging from nominally 3 nm to 13 nm. Upon irradiation with white light at 5 K, all samples exhibit photoinduced magnetism. Magnetization studies indicate that the smallest particles are superparamagnetic, while the largest ones are ferrimagnetic with long-range ordering temperatures ($T_c \sim 17$ K) and coercive fields ($H_c \sim 250$ G) varying with particle size in a manner consistent with the predictions of finite-size scaling.

¹Supported by NSF DMR-0305371 (MWM) and DMR-0543362 (DRT).

D. M. Pajerowski
Department of Physics, University of Florida

Date submitted: 15 Nov 2006

Electronic form version 1.4