

Abstract Submitted
for the MAR07 Meeting of
The American Physical Society

Electron-electron interaction effects on cross-polarized optical absorption in semiconducting single-walled carbon nanotubes (S-SWCNTs)¹ ZHENDONG WANG, SUMIT MAZUMDAR, University of Arizona — Within the tight binding theory of S-SWCNTs optical transitions polarized transverse to the nanotube axis, E_{12} and E_{21} , are degenerate, and occur at $(E_{11} + E_{22})/2$, where E_{11} and E_{22} are the optical transitions polarized along the nanotube axis. Electron-electron interactions split the degeneracy of the transverse transitions, giving new eigenstates that are a redshifted optically forbidden odd superposition and a blueshifted (by several tenths of 1 eV) optically allowed even superposition of the one-electron excitations². Recent experiments³ have confirmed our qualitative prediction. Here we report quantitative calculations of the longitudinal versus transverse optical absorptions in the four S-SWCNTs studied by Miyauchi *et al.*, within a π -electron Hamiltonian with long range Coulomb interactions⁴. We make detailed comparisons between experiments and theory. We also comment on the role of electron hoppings beyond nearest neighbor and the binding energy of the transverse exciton.

¹Supported by NSF-DMR-0406604

²H. Zhao and S. Mazumdar, Phys. Rev. Lett. **93**, 157402 (2004)

³Y. Miyauchi, M. Oba and S. Maruyama, Phys. Rev. B, accepted (2006)

⁴Z. Wang, H. Zhao and S. Mazumdar, Phys. Rev. B **74**, 195406 (2006)

Sumitendra Mazumdar
University of Arizona

Date submitted: 17 Nov 2006

Electronic form version 1.4