

Abstract Submitted  
for the MAR07 Meeting of  
The American Physical Society

**A plausible model for the digital response of p53 to DNA damage**

GUSTAVO STOLOVITZKY, IBM T.J. Watson Research Center, Yorktown Heights, New York, LAN MA, The University of Texas Southwestern Medical Center, Dallas, Texas, JOHN WAGNER, J. JEREMY RICE, IBM T.J. Watson Research Center, Yorktown Heights, New York, HU WENWEI, Cancer Inst of New Jersey, Univ. of Med and Dentistry of NJ, New Brunswick, New Jersey, ARNOLD LEVINE, School of Natural Sciences, Institute for Advanced study, Princeton, New Jersey — The single-cell response of p53 to ionizing radiation (IR) is such that the number of oscillations of p53 shows dependence on the radiation dose. We present a model of this phenomenon. In our model, double strand break (DSB) sites induced by IR interact with a limiting pool of DNA repair proteins, forming complexes that are sensed by ATM, a protein kinase that activates p53 once phosphorylated by DNA damage. The ATM sensing module switches on or off the downstream p53-mdm2 negative feedback loop. Our simulations show that by assuming stochasticity in the initial number of DSBs and the DNA repair process, p53 and Mdm2 exhibit a coordinated oscillatory dynamics upon IR stimulation in single cells, with a stochastic number of oscillations whose mean increases with IR dose.

Gustavo Stolovitzky  
IBM T.J. Watson Research Center, Yorktown Heights, New York

Date submitted: 19 Nov 2006

Electronic form version 1.4