

Abstract Submitted
for the MAR07 Meeting of
The American Physical Society

Temperature and Pump Pulse Dependence of Superfluorescence from InGaAs/GaAs Multiple Quantum Well in High Magnetic Fields¹

XI-AOMING WANG, YOUNG-DAHL JHO, JINHO LEE, DAVID REITZE, University of Florida, JUNICHIRO KONO, Rice University, ALEXEY BELYANIN, VITALY KOCHAROVSKY, Texas A&M University, GLENN SOLOMON, Stanford University, XING WEI, STEPHEN MCGILL, NHMFL — Using intense near-IR ultrashort pulse laser excitation, we investigate the characteristics of cooperative emission (superfluorescence) from dense electron hole magneto-plasmas in InGaAs/GaAs MQW in high magnetic fields as a function of temperature and excitation pulsewidth. We find strong narrow line emissions from 0-0 and 1-1 Landau levels (LLs), with thresholds depending on magnetic field and temperature. Varying the excitation pulsewidth (180 fs – 60 ps) and fluence (0.1 – 1 mJ/cm²), we observe qualitative changes in the emission strengths from different LLs. The strong emissions from 0-0 and 1-1 LL excited with short and long pulses are obviously different. Mechanisms of the temperature and excitation pulse width effect on the strong emission are presented.

¹Supported by the NSF through grant DMR-0325499 and by the NHMFL through an IHRP grant.

David Reitze
University of Florida

Date submitted: 29 Nov 2006

Electronic form version 1.4