

Abstract Submitted  
for the MAR07 Meeting of  
The American Physical Society

**Vibrational Modulation of High Harmonic Generation in SF<sub>6</sub>**<sup>1</sup>

ZACHARY WALTERS, Department of Physics and JILA, University of Colorado, Boulder, Colorado 80309-0440, USA, STEFANO TONZANI, Northwestern University Chemistry Dept. 2145 Sheridan Rd. Evanston, IL 60208-3113, CHRIS H. GREENE, Department of Physics and JILA, University of Colorado, Boulder, Colorado 80309-0440, USA — In a recent experiment performed at JILA (N. Wagner et al, PNAS **103** 13279, 2006), a molecule is hit by two pulses: the first stimulates Raman-active vibrations while the second generates high-order harmonics. The harmonic intensity oscillates as a function of delay time between the two pulses, with oscillation frequencies equal to those of the Raman-active modes. We interpret this oscillation as a form of quantum interference between neighboring vibrational states of the molecule. Nonzero derivatives of the ionization and recombination amplitudes with respect to nuclear coordinates give the molecule some amplitude to change vibrational states during the high harmonic process. We present a theoretical description of vibrational high harmonic modulation and compare with the experimental results of Wagner *et al.*

<sup>1</sup>This work was supported in part by the Department of Energy, Office of Science, and in part by the NSF EUV Engineering Research Center.

Zachary Walters  
Department of Physics and JILA, University of Colorado

Date submitted: 02 Dec 2006

Electronic form version 1.4