

Abstract for an Invited Paper  
for the MAR08 Meeting of  
The American Physical Society

### **Controlling physical parameters of layer-structured nitride-halide superconductors<sup>1</sup>**

YASUJIRO TAGUCHI, RIKEN

Metal-intercalation into band insulators sometimes affords superconductors, well-known examples of which are carbon-based materials, such as fullerenes and graphite. Layer-structured nitride-halide  $\text{Li}_x\text{ZrNCl}$  and  $\text{Li}_xM_y\text{HfNCl}$  ( $M$  denotes molecule) belong to another class of intercalation-induced superconductors with relatively high  $T_c$ , in which doping level and interlayer distance (and hence interlayer hopping interaction) can independently be controlled by changing Li concentration and the size of the co-intercalated molecule. The controllability provides a unique and interesting opportunity to investigate the effect of the two important physical parameters on  $T_c$  in a single system. Recent progress in the synthesis technique enabled us to obtain for the first time a series of single-phase samples of  $\text{Li}_x\text{ZrNCl}$  with finely controlled doping-levels which were notoriously difficult to prepare. Using these samples, we have established[1] an electronic phase diagram to find anomalous doping evolution of  $T_c$ , which takes a maximum value on the verge of superconductor-insulator transition. Based on this phase diagram and the results of systematic Raman scattering and transport measurements, we will discuss possible roles in producing relatively high  $T_c$  played by charge fluctuation and reduced disorder scattering in the layered structure reminiscent of modulation-doped semiconductors. We will also briefly refer to our very recent results on the Hf-based materials in which both of the doping level and interlayer distance were varied.

[1] Y. Taguchi *et al.*, Phys. Rev. Lett. **97**, 107001 (2006)

<sup>1</sup>This work was done in collaboration with A. Kitora, T. Takano, and Y. Iwasa at IMR, Tohoku University.