

Abstract Submitted
for the MAR08 Meeting of
The American Physical Society

Saturated nucleate pool boiling of oxygen under magnetically-enhanced effective gravity¹ T. A. CORCOVILOS², M. E. TURK³, California Institute of Technology - Division of Physics, Math, and Astronomy, Pasadena, CA, USA, D. M. STRAYER, California Institute of Technology - Jet Propulsion Laboratory, Pasadena, CA, USA, N. N. ASPLUND, N.-C. YEH, California Institute of Technology - Division of Physics, Math, and Astronomy, Pasadena, CA, USA — We investigate the effect of enhancing gravity on saturated nucleate pool boiling of oxygen for effective gravities (g_{eff}) of $1g$, $6g$, and $16g$ ($g = 9.8 \text{ ms}^{-2}$) at a saturation pressure of 760 torr and for heat fluxes of $10 \sim 3000 \text{ Wm}^{-2}$. The effective gravity on the oxygen is increased by applying a magnetic body force generated by a superconducting solenoid. We measure the heater temperature (expressed as a reduced superheat) as a function of heat flux and fit this data to a piecewise power-law/linear boiling curve. At low heat flux ($\lesssim 400 \text{ Wm}^{-2}$) the superheat is proportional to the cube root of the heat flux. At higher heat fluxes, the superheat is a linear function of the heat flux. The value of the transition heat flux separating these two regions is proportional to $g_{\text{eff}}^{0.25}$, indicating a possible link to the critical heat flux.

¹Funding by NASA

²Current address: Rice University, Houston, TX, USA

³Current address: Cornell University, Ithica, NY, USA

T. A. Corcovilos
California Institute of Technology

Date submitted: 02 Dec 2007

Electronic form version 1.4