

Abstract Submitted  
for the MAR08 Meeting of  
The American Physical Society

**Search for Magnesium Diboride like Binary Superconductors<sup>1</sup>**

O. PAUL ISIKAKU-IRONKWE, The Center for Superconductivity Technologies[TCST], Abuja FCT, NIGERIA — Efforts to create compounds iso-electronic and iso-structural with magnesium diboride and also superconducting have so far met with limited success. Doping the Mg-site or the B-site have also not yielded higher transition temperatures. They have either been non-superconducting or have lower transition temperatures, Tc. Searching for magnesium diboride-like compounds using the electronegativity of MgB<sub>2</sub> (1.7333) has yielded a rich family of potential superconductors. The search has been facilitated using the recently designed ElectroNegativity Spectrum Maps [ENSMaps] of binary systems A<sub>x</sub>B<sub>y</sub>. Here we display the potential families. Using the relationship between Tc and atomic mass, we estimate the transition temperatures of the most likely MgB<sub>2</sub>-like binary superconductors. We also suggest materials that can be doped to give an electronegativity of 1.7333.

<sup>1</sup>with research support from Mr. Ferguson Uzoma and Institute for Superconducting and Electronic Materials[ISEM], Wollongong NSW Australia

O. Paul Isikaku-Ironkwe  
The Center for Superconductivity Technologies[TCST], Abuja FCT, NIGERIA

Date submitted: 02 Dec 2007

Electronic form version 1.4