

Abstract Submitted
for the MAR08 Meeting of
The American Physical Society

Non-linear conductance of a short quantum point contact¹ TAIMIN LIU, MARYAM TORABI, AMIR MAHARJAN, ANDREI KOGAN, University of Cincinnati, MICHAEL MELLOCH, Purdue University, STEVEN HERBERT, Xavier University — We have measured non-linear conductance G of a very short, less than 80 nm in lithographic length, quantum point contact as a function of the source-drain voltage V_{sd} and gate voltage V_g at the device lattice temperature $T < 20$ mK. The width/length ratio of the QPC is approximately 2.5. We observe several well-resolved plateaus in G at $V_{ds}=0$, but find no prominent zero-bias peak in $G(V_{ds})$ reported by several groups in longer, lower aspect ratio contacts at near-opening gate voltage [1,2]. The peak is believed to arise due to Kondo-like correlations between a quasi-local magnetic state in the constriction [1,3] and the device leads. Our data suggest that the quasi-bound spin state does not form in short QPCs and agree qualitatively with the recent predictions [3]. [1] S.M. Cronenwett et. al. *Physical Review Letters*, 88(226805), 2002. [2] E.J. Koop et. al. *J Supercond Nov Magn*, 20:433, 2007.[3] Tomaz Rejec and Yigal Meir. *Nature*, 442:900, 2006.

¹Supported by the University of Cincinnati start-up funds

Andrei Kogan
University of Cincinnati

Date submitted: 27 Nov 2007

Electronic form version 1.4