

Abstract Submitted
for the MAR08 Meeting of
The American Physical Society

Isotope Effect in High-Tc Superconductors DALE HARSHMAN, Physik Research Corp., JOHN DOW, Arizona State University, ANTHONY FIORY, New Jersey Institute of Technology — For high-Tc superconductors in which transition temperatures, T_c , are reduced by doping, the oxygen isotope effect (OIE) coefficient in T_c is shown to increase systematically with the pair-breaking rate and with the valence difference between the substituted and native ions. Moreover, the OIE tends to zero as one approaches optimum (or ideal) stoichiometry at which the quality of the superconducting condensate is maximized. In materials with isovalent substitutions, e.g., Sr for Ba or Zn for Cu in YBCO, the small OIE of the parent compound is magnified, owing to pair-breaking disorder. In materials with heterovalent substitutions, e.g., La or Pr for Ba, where carrier densities are necessarily changed, pair breaking induces a much larger OIE. A seminal case is Pr-doped YBCO, where the decrease in T_c observed with Pr doping arises from pair-breaking caused by Pr-on-Ba-site defects. Without the defects, T_c is invariant, providing strong evidence against phononic mechanisms. The fact that T_c drops when Pr substitutes for Ba, but not for Y, indicates that the superconducting hole condensate resides in the BaO layers, where pair-breaking degrades T_c and dramatically increases the OIE. Superconductive pairing modeled on Coulomb coupling between the hole and the electron layers is shown to resolve the shortcomings in electron-phonon interactions.

Anthony Fiory
New Jersey Institute of Technology

Date submitted: 27 Nov 2007

Electronic form version 1.4